

Mathematical Fuzzy Logic

Ph.D. course

Tommaso Flaminio and Carles Noguera

Department of Mathematics and Computer Science

University of Siena

May – June 2009

Aims of the course: To introduce the basic notions of Mathematical Fuzzy Logic and some of the open problems in its agenda.

Topics:

1. Introduction to the algebraic study of propositional logics. The cases of intuitionistic and classical logic. The general notion of algebraizable logic.
2. Gödel-Dummett logic as an extension of intuitionistic logic. Product and Lukasiewicz logic. Algebraic semantics. Standard semantics given by continuous t-norms. Finite strong standard completeness properties and failure of strong standard completeness.
3. Hájek's BL logic as the logic of all continuous t-norms and their residua. Algebraic semantics, ordinal sums and standard completeness properties. Hilbert style calculus.
4. Residuation and left-continuity. MTL as the logic of all left-continuous t-norms and their residua. Algebraic semantics, Hilbert style calculus and Local Deduction Theorem. Real embedding property.
5. Core fuzzy logics. Completeness properties with respect to distinguished semantics (real, rational, hyperreal and finite chains): methods and equivalencies.
6. First-order predicate fuzzy logics: axiomatization and semantics. Completeness properties.
7. Modal fuzzy logics.

References

- [1] P. AGLIANÒ AND F. MONTAGNA. Varieties of BL-algebras I: general properties, *Journal of Pure and Applied Algebra*, 181 (2003) 105–129.

- [2] W. J. BLOK AND D. PIGOZZI. Algebraizable logics, *Memoirs of the American Mathematical Society* 396, vol 77, 1989.
- [3] S. BURRIS AND H. P. SANKAPPANAVAR. *A course in Universal Algebra*, Springer Verlag, New York, 1981.
- [4] R. CIGNOLI, F. ESTEVA, L. GODO AND A. TORRENS. Basic Fuzzy Logic is the logic of continuous t-norms and their residua, *Soft Computing* 4 (2000)
- [5] P. CINTULA, F. ESTEVA, J. GISPERT, L. GODO, F. MONTAGNA AND C. NOGUERA. Distinguished algebraic semantics for t-norm based fuzzy logics: methods and algebraic equivalencies, *Annals of Pure and Applied Logic*, 2009, in press.
- [6] M. DUMMETT. A propositional calculus with denumerable matrix, *The Journal of Symbolic Logic* 24 (1959) 97–106.
- [7] F. ESTEVA, J. GISPERT, L. GODO AND F. MONTAGNA. On the standard and Rational Completeness of some axiomatic extensions of Monoidal t-norm Based Logic, *Studia Logica* 71 (2002) 199–226.
- [8] F. ESTEVA AND L. GODO. Monoidal t-norm based logic: Towards a logic for left-continuous t-norms. *Fuzzy Sets and Systems*, 124: 271–288, 2001.
- [9] T. FLAMINIO. Strong non-standard completeness for fuzzy logics, *Soft Computing* 12 (2008) 321–333.
- [10] T. FLAMINIO AND E. MARCHIONI. Continuous t-norm predicate logics are complete with respect to rational-safe models, Submitted, 2007.
- [11] J. M. FONT, R. JANSANA AND D. PIGOZZI. A Survey of Abstract Algebraic Logic, *Studia Logica* 74 (2003) 13–97.
- [12] P. HÁJEK. *Metamathematics of Fuzzy Logic*, volume 4 of *Trends in Logic-Studia Logica Library*. Dordrecht/Boston/London, 1998.
- [13] P. HÁJEK AND P. CINTULA. On theories and models in fuzzy predicate logics, *The Journal of Symbolic Logic* 71: 863–880, 2006.
- [14] P. HÁJEK, L. GODO AND F. ESTEVA. A complete many-valued logic with product-conjunction, *Archive for Mathematical Logic* 35 (1996) 191–208.
- [15] R. HORČÍK. Standard Completeness Theorem for IIMTL, *Archive for Mathematical Logic* 44 (2005) 413–424.
- [16] S. JENEI AND F. MONTAGNA. A proof of standard completeness for Esteva and Godo’s logic MTL. *Studia Logica*, 70:183–192, 2002.
- [17] P. S. MOSTERT AND A. L. SHIELDS. On the structure of semigroups on a compact manifold with boundary, *Annals of Math.* 65 (1957) 117–143.
- [18] C. NOGUERA. *Algebraic study of axiomatic extensions of triangular norm based fuzzy logics*, Monografies de l’Institut d’Investigació en Intel·ligència Artificial vol. 27, Barcelona, 2007.

[19] L. A. ZADEH. Fuzzy sets, *Information Control* 8 (1965) 338–353.